



Mississippi Corn Promotion Board 2020 Progress Report

Project Title: Trapping Program to Monitor Southwestern Corn Borer Populations in Mississippi

PI: Jeff Gore, W. Crow, A. Catchot, D. Cook, and F. Musser

Department: Delta Research and Extension Center and Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology

Project Summary (Issue/Response)

The southwestern corn borer is one of the most important insect pests of non-Bt corn in Mississippi. Currently, growers are required to plant a refuge of non-Bt corn to satisfy the resistance management plan for Bt corn hybrids. Those refuges are subject to potential yield losses from southwestern corn borer annually. Bt corn hybrids have been widely utilized in Mississippi because they provide excellent control and monitoring for southwestern corn borer in non-Bt corn can be very difficult. Because an economically damaging infestation is difficult to detect, a comprehensive trapping program is needed in Mississippi to help growers properly time insecticide applications in the non-Bt refuge and minimize losses from this insect.

To monitor southwestern corn borer populations throughout the year, pheromone traps were placed at various locations throughout Mississippi. Every attempt was made to locate the traps adjacent to a non-Bt corn field. A total of 64 locations were monitored throughout the state that included all of the major corn producing regions. The traps were identified by county and nearest town for reporting purposes. The traps were monitored weekly and the number of southwestern corn borers per trap were recorded. Trap counts were reported weekly on the Mississippi Crop Situation blog (www.mississippi-crops.com) and through other means to ensure that the information was widely disseminated to growers, consultants, and field scouts.

Project Results/Outcomes

A total of 84 southwestern corn borer trapping locations were monitored from May through August in 2020. Similar to previous years, trap catches were highly variable from trap to trap within a region. Overall populations were considerably higher and persisted longer in 2020 than in previous years. Based on trap catches, the first generation of moths peaked from late-May to early-June. Populations of the second generation began increasing the last week of June and peaked during the third week of July. However, populations in individual traps persisted throughout July and most of August. Individual traps across the state exceeded the current threshold from July 4 to August 8 in 2020. This is much different than what has been observed in previous years. Overall, the corn crop was later than in previous years, so more acres of corn were at a susceptible stage when populations peaked, especially in the north Delta. The highest trap counts were recorded from the central to northern part of the state (North of Hwy 82). Similar to previous years, the highest trap counts occurred in Coahoma, Leflore, and Tate counties during 2020. Additionally, several traps across the state that typically trap low corn borer numbers experienced higher than normal corn borer populations. Higher than normal populations were detected in Carroll, Sunflower, Humphreys, and Quitman counties during 2020. Based on the high level of variability from trap to trap in this survey, it is highly recommended that growers and consultants utilize pheromone traps to monitor southwestern corn borer in individual fields. Although trapping is recommended in individual fields, these surveys provide tremendous value to corn growers in Mississippi.




Project Results

They provide an initial indication of when southwestern corn borer populations are beginning to increase across the state, and signal the timings when scouting should be intensified in individual fields. Additionally, these surveys can be used to trigger insecticide applications in fields where traps are not being used by consultants and growers. Results of these surveys were reported weekly on the Mississippi Crop Situation blog and communicated through phone conversations and text messages to growers, consultants, and retail scouts.



Southwestern Corn Borer Trapping
Mississippi - 2020



Mean and maximum number of southwestern corn borers caught in traps across Mississippi during 2018. The maximum represents the highest number of moths that were captured in any one trap during a particular week.

Project Impacts/Benefits

The results of these trapping efforts are an important first step in determining when damaging levels of southwestern corn borer are likely to occur in Mississippi. Because field scouting for infestations of this pest is very difficult and because timing of foliar applications of insecticides is critical for effective control, these results are important for helping field scouts, consultants and growers determine when to spray. As the percentage of non-Bt corn acres increases due to low commodity prices, these efforts will become more important to ensure effective control and economical production of field corn in Mississippi.

Project Deliverables

The results of this survey were presented weekly in the Mississippi Crop Situation Newsletter (www.mississippi-crops.com) and through personal phone calls from numerous field scouts and consultants throughout the state.

Results have also been presented at numerous field days, grower meetings, and workshops throughout the state.